



Our new line of products with all the experience of and NUTRITION

Trust the specialists
www.andnutrition.com
Lorca, 27. 28231 Las Rozas, Madrid (Spain)



and AMINOVIT

Pharmaceutical form:

Oral solution

Target species:

Bovine – Goats – Ovine – Porcine – Poultry - Rabbits

Presentation:

250 ml, 1L and 5L containers (with dosifier).

Composition: (in 1 litre)

Folic acid, 250 mg	Niacin, 25 g
L-Alanine, 6,35 g	Sorbitol, 105 g
L-Arginine, 9,9 g	DL-Carnitine, 50 g
Aspartic acid, 9,35 g	L-Threonine 3,48 g
Biotin, 20 mg	L-Tryptophan, 2 g
L-Cystine, 1,70 g	Vitamin B1, 2,5 g
Choline chloride, 28 g	Vitamin B2, 1,75 g
Pantothenic acid, 7 g	Vitamin B6, 1,1 g
L-Phenylalanine, 8,35 g	Vitamin B12, 1,25 mg
Glycine, 13,55 g	Vitamin A, 875.000 U.I
Glutamic Acid, 7,55 g	Vitamin D3, 66.600 U.I
L-Histidine, 7,70 g	Vitamin E, 2 g
L-Isoleucine, 4,8 g	Vitamin K3, 2 g
L-Leucine, 16,6 g	
L-Lysine, 40 g	
DL-Methionine 10 g	

Indications:

Appetite stimulant, energetic contribution. It regulates hepatic and kidney-related functions.

Dosage:

Oral administration. General dose: 1 ml / 1 litre of water (1 l / 1000 l)
Adult bovine animals: 40 ml/day during 10 days. Calves: 30 ml/day during 10 days. Reproductive pigs: 30 ml/day during 10 days. Growing pigs: 2 ml/day during 10 days. Birds: 4 ml/Kg fodder during 5 days. Rabbits, partridges and quails: 1 ml/Kg fodder during 10 days. Sheep and goats: 6 ml/day during 10 days. Lambs and young goats: 2 ml/day during 10 days.



and SELENVIT

Pharmaceutical form:
Oral liquid

Target species:
Goats – Ovine – Porcine – Broilers - Layers

Presentation:
1 and 5 Litres containers.

Composition: per 100 ml
Propylene glycol
Vitamin A, 135.000 UI
Vitamin E, 5.000 mg
Selenium (Sodium selenite), 23 mg
Citric acid, 500 mg

Indications:
Critical period of growth (weaning, transport, stress, summer ...)

Dosage:
Oral administration.
As a general dose: use 0,5 to 1 ml / 1 litre of water (500 ml to 1 l. per 1000 l.)
Lambs, young goats and piglets: 1 ml/animal. Renew in 15 days time.
Pay attention to the dosage due to the toxicity of selenium.



and LIVERPROTECT

Pharmaceutical form:

Oral dust

Target species:

Bovine – Goats – Sheep – Porcine - Poultry

Presentation:

1,5 and 25 Kg containers.

Composition:

Vitamin A, 800.000 U.I

Vitamin D3, 160.000 U.I.

Vitamin E, 2g

Vitamin C, 9 g

Vitamin B1, 500 mg

Vitamin B2, 1 g

Vitamin B6, 500 mg

Vitamin B12, 12 mcg

Choline, 17 g

Sorbitol, 160 g

Betaine, 25 mg

DL-Methionine, 40 g

Indications:

It is a hepatic protector. It stimulates the appetite and it avoids the loss of weight in high productive periods. It prevents stress.

Dosage:

Oral administration.

Birds: 2-4 g/Kg of fodder.

Pigs: 10 g/Kg of fodder.

Calves: 30 g/animal/day

Dairy cows: 40 g/animal/day at least during 28 days.

Ovine and goats animals: 5 g/animal/day during 21-28 days just before the parturition.

Fighting bulls: 70 g/bull/day during 7 days.

Oxen: 50 g/animal during 1 month.



and HIDRATION LQ

Pharmaceutical form:
Oral solution

Target species:
Bovine – Goats – Horses – Sheep – Porcine - Poultry

Presentation:
1, 5 and 25 L containers

Composition:
Glucose monohydrate, 27,8 g
Sodium chloride, 5,33 g
Glycine, 3,85 g
Potassium phosphate monobasic, 2,55 g

Indications:
It stabilizes the hydric and electrolyte balance. It contributes to the animal's convalescence after diarrhoeic processes.

Dosage:
Oral administration. General dose: 50 – 75 ml per litre.
Lambs and young goats: 75 ml diluted in 1 L. of drinking water. Administer 500 ml to 1 L distributed in several doses.
Dehydrated calves: 75 ml diluted in 1 L of drinking water. Administer 2 L distributed in several doses during the same day. It can be supplied as one pleases.
Piglets to be taken from the nipple suffering from diarrhoea: 75 ml diluted in 1 L of drinking water. Administer 500 ml to 1 L distributed in several doses.
Colts: 75 ml diluted in 1 L of drinking water. Administer 2 L to 3 L distributed in several doses. Recommended period of use: 1 to 7 days.



and HIDRATION PD

Pharmaceutical form:
Soluble powder

Target species:
Bovine – Goats – Horses – Sheep – Porcine - Poultry

Presentation:
5 kg containers and 25 Kg bags

Composition:
Per Kg.:
Sucrose, 425 g
Sodium chloride, 80 g
Calcium chloride, 2 g
Potassium chloride, 5 g
Sodium hydrogen carbonate (sodium bicarbonate), 40 g
Citric acid, 5 g
Glycine, 1 g

Indications:
It stabilizes the hydric and electrolyte balance. It contributes to the animal's convalescence after diarrhoeic processes.

Dosage:
Oral administration. 10 – 25 grams per litre of warm water. Repeat the dose in 12 hours time.

Transport: 50 g before starting the trip. Repeat the dose when arriving in your destination. Recommended treatment: 1 to 7 days (or 1 day after stop of symptoms).



and HIDRATION GEL

Pharmaceutical form:
Oral gel

Target species:
Bovine – Sheep – Goats - Pigs

Presentation:
Box with a 15 g dosifier syringe.

Composition:
Dextrose
Sodium chloride
Calcium chloride
Magnesium chloride
Potassium sorbate
By-products of plants

Indications:
It stabilizes the hydric and electrolyte balance. It is suited to the prevention and treatment of disorders such as diarrhoea and for convalescence processes.

Dosage:
Oral administration. Lambs and young goats: 5 g/animal/day.
Calves: 15 g/animal/day. Recommended period of use: 1 to 7 days (1 to 3 days if it is the only aliment).



and BABY POWER

Pharmaceutical form:
Liquid

Target species:
Bovine – Sheep – Goats – Cats – Dogs – Horses – Ovine – Porcine - Rabbits

Presentation:
250 ml container with 2ml pump dosifier (125 doses).

Composition:
Iron (Iron lactate trihydrate), 94,7 mg
Dextrose monohydrate, 50 mg
Folic acid, 13 mcg
Vitamin A, 16,90 UI
Vitamin D3, 1,30 UI
Vitamin E, 0,073 mg
Vitamin B1, 5 mcg
Vitamin B2, 8 mcg
Vitamin B6, 8 mcg
Nicotinic acid, 14 mcg
Vitamin C, 0,2 mg

Indications:
An extraordinary contribution of Iron, vitamins and gluco-energetic substances that allow us to provide the animals with the essential elements they need in order to get the maximum equilibrium in their organisms and stimulate their immune systems by a progressive effect.

Dosage:
Oral administration.
Piglets: 1-2 ml (1 or 2 pulses) in their 3 first days of life.

Properties:
It gives energy, iron and vitamins to the new born animal and improves its immunity



MEGAMACHO

Pharmaceutical form:

Soluble powder

Target species:

Bovine – Goats – Horses – Sheep – Porcine (sows and boars) – Poultry. Specially designed to improve reproduction in boars, gilts, sows, cows, bulls

Presentation:

5 kg

Composition:

Calcium carbonate, monocalcium phosphate, sodium chlorine (sea salt) magnesium oxide. Vitamins. Minerals. Amino acids

Calcium: 24 %

Phosphorus: 5,4 %

Sodium 1,2 %

Magnesium: 0,6 %

Suffer: 0,04 %

Vitamin A (E672). 67.500 UI	Iron (E1) (sulfato monohidrato) .400 mg
Vitamin D ₃ (E671). 13.500 UI	Iodine (E2) (Ioduro potásico) . 100 mg
Vitamin B ₁ . 100 mg	Cobalt (E3)(sulfato heptahidratado) . 100 mg
Vitamin B ₂ . 200 mg	Copper(E4) (sulfato pentahidrato) . 300 mg
Vitamin B ₆ . 10 mg	Manganesum (E5) (sulfato monohidrato)640 mg
Vitamin B ₁₂ . 15 mg	Zinc (E6) (oxido) 720 mg
Vitamin C. 100 mg	L-Lisine 1.950 mg
Vitamin E (alfa tocoferol). 75 mg	DL-Metionine 10.000 mg
Vitamin K ₃ . 25 mg	DL-Carnitine HCl 10.000 mg
Nicotinic acid: 1.250 mg	Mixture of flavours: 5.000 mg
Calcium pantotenate 600 mg	Anticaking agents:
Cholin clorure 25.000 mg	Sepiolite (E562) 166.000 mg

Dosage:

Cows and bulls: 100 g/animal/day; Sheep and goats: 10 g/animal/day; Reproductive sows and boars: 40 g/animal/day; Growers and fatteners: 2 g/10 Kg.weight.

Poultry: 1 g/animal/day



VIRUBAC

Pharmaceutical form:
Liquid

Target species:
Installations. Buildings of all animal species.

Presentation:
1, 5 and 25 L containers.

Composition:
Glutaraldehyde 14%
Didecyl Dimethyl Ammonium Chloride, 10%
Surface- actives and excipients c.s.p. 100%

Dosage: (from 0,33% to 1%)
Porcine animals:

Maternity: general weekly disinfection 1 L/300 L of water.
Sanitary vacuum: 1 L/100 L of water.
Breeding place: Regular disinfection: 1 L/200 L of water.
Disinfection by pathology: 1 L/100 L of water.
Medicinal foot bath: 1 L/100 L of water.

Aviculture and rabbits breeding:

Fattening and over-breeding: Sanitary vacuum: 1 L/200 L of water.
Grounds: 1 L/50 L of water.
Egg- laying and bacteria: regular disinfection: 1 L/200 L of water.
Disinfection with pathologies: 1 L/100 L of water.
Cages and material: 1 L/200 L of water.

Bovine, sheep, goats and equine animals:

Stables: Habitual weekly disinfection: 1 L/200 L of water.
Disinfection with pathologies: 1 L/100 L of water.
Medical foot bath: 1 L/100 L of water.

Installations, Material and Equipment: 1 L/150 L of water.

Drinking water circuits: keep during one hour the product dissolved in 1 L/150 L of water in the piping circuits, eliminate it afterwards and then clean it with water.

Animal's transport: General disinfection: 1 L/150 L of water.

Properties:

Virubac incorporates two active principles widely used, whose synergic action allows a complete bacterial and fungicide disinfection.

Trust the specialists
www.andnutrition.com
Lorca, 27. 28231 Las Rozas, Madrid (Spain)



ALUDERMIN

Pharmaceutical form:
Topic aerosol

Target species:
Bovine – Sheep – Goats – Cats – Dogs – Horses – Ovine – Porcine - Rabbits

Presentation:
270 ml containers and 120 g (net quantity). Spray.

Composition:
Micronized aluminium, 50 %
Excipient c.s.

Indications:
Protection of wounds against dirtiness, insects, etc.

Dosage:
After having cleaned and dried the skin, apply a small quantity of the product over the skin surface to be protected.
Keep the spray at a distance of 20 centimetres.